

שם: פרופ' משה (מושיק) ענבר

שם המוסד: אוניברסיטת חיפה

דוא"ל: minbar@research.haifa.ac.il

תחומי עניין ומחקר: אינטראקציות בין בעלי חיים לצמחים
רשימת מקוצרת של פרסומים מדעיים רלבנטיים 'לכלנית'

Inbar M, Eshel A & Wool D (1995) Interspecific competition among phloem – feeding insects mediated by induced host plant sinks. *Ecology*. 76:1506-1515

Inbar M & Wool D (1995) Phloem-feeding specialists sharing a host tree: Resource partitioning minimizes interference competition among galling aphid species. *Oikos*. 73:109-119

Wool D, Manheim O & Inbar M (1997) Return flight of sexuparae of galling aphids to their primary host trees: implications for differential herbivory and gall (Aphidoidea: Pemphigidae, Fordinae) abundance. *Annals of the Entomological Society of America*. 90:341-350

Wool D & Inbar M (1998) Colonization of ecological islands: galling aphid populations on recovering *Pistacia* trees after destruction by fire. *European Journal of Entomology*. 95:41-54

Inbar M, Doostdar H, Sonoda RM, Leibe GL & Mayer RT (1998) Elicitors of plant defensive systems reduce insect densities and disease incidents. *Journal of Chemical Ecology*. 24:135-149

Inbar M (1998) Competition, territoriality and maternal defense in a gall-forming aphid. *Ethology, Ecology and Evolution*. 10:159-170

Inbar M, Doostdar H & Mayer RT (1999) The effects of sessile whitefly nymphs (*Homoptera: Aleyrodidae*) on leaf chewing larvae (*Lepidoptera: Noctuidae*). *Environmental Entomology*. 28:353-357

Inbar M, Doostdar H, Leibe GL & Mayr RT (1999) The role of plant rapidly induced responses in asymmetric interspecific interactions among insect herbivores. *Journal of Chemical Ecology*. 25:1961-1979

Inbar M, Doostdar H & Mayer RT (2001) Suitability of stressed and vigorous plants to various insect herbivores. *Oikos*. 94:228-235

Lev-Yadun S, Dafni A, Inbar M, Izhaki I & Ne'eman G (2002) Colour patterns in vegetative parts of plants deserve more research attention. *Trends in Plant Science*. 7:59-60

Auslander M, Nevo E & Inbar M (2003) The effects of slope orientation on plant growth, developmental instability and susceptibility to herbivores. *Journal of Arid Environments*. 55:405-416

Inbar M, Wink M & Wool D (2004) The evolution of host plant manipulation by insects: Molecular and ecological evidence from gall-forming aphids on *Pistacia*. *Molecular Phylogenetics and Evolution*. 32:504-511

- Adar E, Inbar M, Gal S, Gan-Mor S & Palevsky E (2014) Pollen on-twine for food provisioning and oviposition of predatory mites in protected crops. *BioControl*. 59:307-317
- Rand K, Bar E, Ben-Ari M, Lewinsohn E & Inbar M (2014) The mono - and sesquiterpene content of aphid-induced galls on *Pistacia palaestina* is not a simple reflection of their composition in intact leaves. *Journal of Chemical Ecology*. 40:632-642
- Kurzfeld-Zexer L, Lev-Yadum S & Inbar M (2015) One aphid species induces three gall types on a single plant: comparative histology of one genotype and multiple extended phenotypes. *Flora*. 210:19-30
- Gish M, Ben-Ari M & Inbar M (2017) Direct consumptive interactions between mammalian herbivores and plant-dwelling invertebrates: prevalence, significance and prospectus. *Oecologia*. 183:347-352
- Rand K, Bar E, Ben-Ari M, Davidovich-Rikanati R, Dudareva N, Inbar M & Lewinsohn E (2017) Differences in monoterpene biosynthesis and accumulation in *Pistacia palaestina* leaves and aphid- induced galls. *Journal of Chemical Ecology*. 43:143-152
- Berman TS, Ben-Ari M, Henkin Z & Inbar M (2018) Immediate and long-term facilitative effects of cattle grazing on a polyphagous caterpillar. *Agriculture, Ecosystems and Environment*. 261:45-53
- Ben-Ari M, Outreman Y, Denis G, Le Gallic JF, Inbar M, Simon JC (2019) Differences in escape behaviour between pea aphid biotypes reflect their host plants' palatability to mammalian herbivores. *Basic and Applied Ecology*. 34:108-117
- Schlesinger D, Davidovich Rikanati R, Volis S, Faigenboim A, Vendramine V, Cattonaro F, Hooper M, Oren E, Taylor M, Sitrit Y, Inbar M, Lewinsohn E (2019) Chemodiversity in *Mandragora* spp. is associated with loss-of-functionality of MoH6H, a hyoscyamine 6b-hydroxylase gene. *Plant Science* 283:301-310